

, Im

# \*\*\*\*\*\*\*\*

The mediceine di move belle for 9 out of every 10 woman dem wey dem tak'am.

#### IN CASE THE MEDICINE No Work?



Make e take the medicine again (12 tablets, 4 tablets under your tongue until them melt completely, every 3, 3 hours) or make e go go get Manual Vacuum Aspiration (MVA).-for this method, dem go insert something like syringe inside your woman skin for suck out any part of the belle wey e remain or no commot. But if the belle no commot after vou don take the MISO, small risk dey say the pikin wey you go born fit get problem.

#### **THE BELLE COMMOT ALL?**

Even if the woman feel say e no dey with belle again, e be important say me e confirm say the belle don commot completely. E go either do scan, or pregnancy test 3-4 weeks after e don move the belle.

in case the **woman no bleed**, **or e bleed only small** (wey blood no commot like menstruation and e no get blood clot) and sign dem for pregnancy still continue after e don take the tablets dem, she fit like still get belle. Make e hurry go do scan for confirm if the belle still dey or e don commot. If e no fit do the scan, e fit take the tablets them again!

# AFTER THE BELLE DON COMMOT

After the belle don commot, the woman fit expect for continue for bleed for about 1-3 weeks; but no be the same for all woman them. After everything, the woman get for confirm say the belle don commot completely.

We di dvise say make woman dem no put anything inside their woman skin, wash or sleep with man pikin until that plenty blood don stop for commot; about 2 days after them don use MISO for move the belle.

Woman fit get belle immediately after e don move another belle. If e no want get belle immediately, e must start for di do thing dem wey them no go mke e get another belle wey e want keep'am- like for use condom, and so on.

### CONTACTS

Women HelpWomen WEB: HTTPS://WOMENHELP.ORG/ EMAIL: INFO@WOMENHELP.ORG TWITTER: @WOMENHELPORG FACEBOOK: WOMENHELPWOMENINTERNATIONAL

MAMA Network: Mobilizing Activists around Medical Abortion (MAMA). WEB: WWW.MAMANETWORK.ORG



# WETTI BE MISOPROSTOL?

Na medicine wey dem di use'am for all side for this world for move belle wev vou no wan'am and for cure gastric ulcers. For inside medicine store, dem di sellam with different different kind name dem. Like cytotec, isovent, misoclear, misoFem, cyrux of tomispral. For plenty African country dem, them diuse this medicine for prevent and for stop blood after woman don born.



For move belle with no problem.

For finish for move belle wey dem no do'am fine. For prevent and stop blood after woman don born. Them fit us'am too for treat gastric ulcer.

#### HOW E DI WORK?

Misoprostol go mak'am make e be like say you don get miscarriage. E go make that road we pikin di pass soft and open we e go easy for the belle for commot.

#### WHEN DEM FIT USAM For move belle with No problem?




You fit us'am within the first 12 weeks weh you de with belle starting

from the first day after

your last for see your


period.



The pains and<br/>cramps/pains go<br/>plenty pass as e di<br/>use for be during<br/>period.80% of the time,<br/>belle di commot<br/>before 24 hours<br/>want finish.

ne, The more the belle ot dons stay, the rs more the blood and cramps/pains fit be.

You fit take pain killers the same time wey you di take the first dose of miso. **Ibuprofen** na the best. But you fit also take **paracetamol**/ **acetaminophen**.



You go get High temperature/fever wey pass 39 degree, for more than 1 day.

WHETTI FIT HAPPEN

The woman fit notice this kind signs the m within

vomit, purge, fever (wey u no di shiver), headache,

the first 24 hours: E neck go turn (nausea), e fit

body di weak. All these one dem na side-effects



39°

>>

24

Plenty pain wey pain killers no fit cool'am and e go continue for few days after you don take the MISO.



Bad smell and discharge go di commot for your woman skin.

## IF YOU GET ANY PALAVA



Go hospital or heakth center immediately.

Workers for hospital and health center get obligation for attend to the woman immediately.

For hospital, dem go give e the same treatment wey them di give woman wey get complications after e don miscarry. **E no be possible make dem know for hospital wether the woman don miscarry or e don try for move belle with MISO.** So woman we y get complication after e don take MISO fit tlk for hospital say don miscarry.

Ð

 $( \bullet )$